


## Drawing up Evogam® Subcutaneous immunoglobulin (SCIg) therapy

1. Wash your hands thoroughly with soap and water and dry with a clean towel.
2. Remove the protective cap from the vial and wipe the rubber stopper with an alcohol wipe.



3. Remove the top cover of the vial adaptor, leaving the adaptor inside the blister pack.




4. Place the vial adaptor over the top of the vial.
5. Remove the outer package of the vial adaptor and discard.



**Use the blister pack to handle the vial adaptor. Take care not to touch the exposed end of the vial adaptor**

6. Pull back on the plunger to fill the syringe with air. The amount of air should be equal to the volume of Evogam® you are about to draw up from the vial (e.g. if you are about to draw up 5ml of Evogam® from a vial, then fill the syringe with 5ml of air).



Remember, take care not to touch the exposed ends of the syringe.

7. Attach the syringe to the vial adaptor by twisting the syringe onto the vial adaptor connection. Gently inject the air into the vial.  
**Turn the vial upside down and then** release the pressure on the plunger and slowly pull back on the plunger to fill the syringe with the Evogam®



8. Disconnect the syringe from the vial adaptor with a twist, leaving the vial adaptor attached to the vial.



9. Repeat for each vial in the dose. Remember to draw back air into the syringe and to gently inject the air into each vial prior to drawing up the solution. **Always use a new vial adaptor for each vial.**

© ASCIA 2017

ASCIA is the peak professional body of clinical immunology and allergy specialists in Australia and New Zealand

**Disclaimer**

This document has been adapted with permission from resources developed by Princess Margaret Hospital Immunology Department (Department of Health, Western Australia). It has been peer reviewed by ASCIA members and is based on expert opinion and the available published literature at the time of review. Information contained in this document is not intended to replace medical advice and any questions regarding a medical diagnosis or treatment should be directed to a medical practitioner. Development of this document is not influenced by commercial organisations.